

# Fulfilling commitments under the ICPD Programme of Action and the Millennium Development Goals

## Advancing women's access to safe abortion


The ICPD Programme of Action (POA) called attention to the health consequences for women of unsafe abortion and called for actions to address this critical public health issue (United Nations 1994).

- Paragraph 7.6 states that abortion care should be an integral part of primary health care.
- Paragraph 8.25 declares that “In circumstances where abortion is not against the law, such abortion should be safe.”
- In 1999, the United Nations’ five-year review of POA implementation strengthened this call to action: “In circumstances where abortion is not against the law, health systems should train and equip health-service providers and should take other measures to ensure that such abortion is safe and accessible.” (United Nations 1999)

Millennium Development Goal (MDG) 5, “Improve maternal health,” seeks a 75 percent reduction of maternal mortality between 1990 and 2015 and includes a target of achieving universal access to reproductive health care. MDG 5 can be met only if unsafe abortion is effectively addressed.

The enormous disparity in maternal mortality and morbidity from unsafe abortion between rich and poor countries underscores that access to safe, comprehensive reproductive health services, including abortion-related care, is a critical development issue. Unsafe abortion constitutes a public health crisis, a social injustice, and a violation of women’s human rights and dignity. Governments worldwide urgently need to adopt policies and interventions to eliminate unsafe abortion as an essential step toward meeting their obligations under the ICPD Programme of Action and achieving MDG 5.

*Estimated annual maternal deaths due to unsafe abortions per 100,000 live births by subregion, 2003.  
Adapted from: WHO 2007*



## A major, preventable cause of maternal deaths and injuries


Unsafe abortion accounts for approximately 13 percent of global deaths from complications of pregnancy and childbirth, and a much higher proportion in many developing countries (WHO 2007). It is one of the easiest causes of maternal mortality to address, through improved access to family planning information and services, high-quality postabortion care, and safe, legal abortion. Yet, unsafe abortion and its root causes remain largely neglected.

Research shows that women have abortions just as frequently where abortion is legally restricted as where it is broadly permitted, but deaths and injuries from unsafe abortion occur at much higher rates in restrictive settings. Around the world, at least 66,500 women die each year from unsafe abortion. In just the past decade, more than half a million women have lost their lives because they lacked access to safe abortion services – women in the prime of life and who most often had children and families to care for. Globally, adolescents and women younger than 24 years account for almost 46 percent of deaths related to unsafe abortion (WHO 2007). The toll of unsafe abortion is especially high in sub-Saharan Africa, where abortion laws generally are very restrictive.

Unsafe abortion also commonly results in long-term and chronic health problems, including reproductive tract infections and infertility. As with deaths related to unsafe abortion, the women most at risk of suffering serious complications are young, poor and reside in rural areas of countries with restrictive abortion laws.

### Percentage distributions of estimated unsafe abortions and associated mortality by age, all developing regions, 2003

Source: WHO 2007



## Progress addressing unsafe abortion since ICPD

Progress in addressing issues underlying abortion-related morbidity and mortality in the last several decades includes a dramatic overall increase in use of modern family planning methods. Between 1960 and 2000, the proportion of married women in developing countries using contraception to prevent unintended pregnancy rose from less than 10 percent to 60 percent (Cleland et al. 2006). Such improvements have been uneven, however, with many poor countries still reporting very low contraceptive prevalence rates.

With support from a range of donors and strong government commitments in many countries, one important achievement since the ICPD has been expanding women's access to high-quality postabortion care – treatment of abortion complications linked to postabortion family planning and other critical reproductive health services. Improving the availability and quality of postabortion care has saved women's lives, improved their health and benefitted health systems by reducing costs and freeing up resources for other needs. But postabortion care alone cannot eliminate deaths and injuries from unsafe abortion.

The 15 years since the ICPD have also brought some improvements in access to safe abortion to the extent allowed by law. In 2003, the World Health Organization (WHO) issued *Safe abortion: technical and policy guidance for health systems*, a groundbreaking resource that paved the way for a number of countries to develop and disseminate national standards and guidelines on the scope and quality of abortion care (WHO 2003). There has also been notable, though still insufficient, progress in training physicians, midwives, and other health-care providers and in making technologies for safe abortion available and affordable, including vacuum aspiration instruments and pharmacological agents as recommended by WHO.

Other organizations and medical groups have issued clinical and ethical guidance on abortion, including the International Planned Parenthood Federation (IPPF 2006) and the International Federation of Gynecology and Obstetrics (FIGO 2006). FIGO's guidelines state that women have the right to access legal, safe, effective, acceptable and affordable methods of contraception and safe abortion services.

Among the most important actions to reduce deaths and injuries from unsafe abortion is liberalizing abortion laws, which about two dozen countries did between 1995 and 2008 (Boland and Katzive 2008, Center for Reproductive Rights 2007). In addition, a number of national, state and provincial governments have already amended or are now considering amending their laws to increase women's access to legal termination of pregnancy. Progress has also been made by



applying the provisions of international human rights treaties to ensure that women are able to exercise their right to terminate a pregnancy safely, based on their rights to health care, privacy and confidentiality, the benefits of scientific progress, and freedom from discrimination, among others.

### **Actions required**

Despite these improvements, much remains to be done if we are to stop women and girls from dying from unsafe abortions. Urgent needs include:

- Ensuring that women, especially those who are most vulnerable, have access in their communities to the full range of reproductive health care, including effective and affordable methods of contraception, postabortion care, and safe abortion, in accordance with their human rights.
- Efforts through community-based approaches and the media to help women understand how to prevent unintended pregnancy, their legal rights to safe abortion, the dangers of unsafe abortion and where to obtain safe abortion care.
- Training more health-care providers, especially midlevel health professionals such as nurses and midwives, in clinical procedures and counseling for abortion care.
- Eliminating violence against women and ensuring that women who are survivors of sexual violence have immediate access to psychological support, treatment of their injuries, emergency contraception, postexposure prophylaxis of HIV, treatment of sexually transmitted infections, and safe, legal abortion.
- Promoting constructive dialogues at the national and local levels about unwanted pregnancy and unsafe abortion.
- Reforming policies and laws to increase access to safe abortion in accordance with human rights principles.

Effectively addressing unsafe abortion as a critical public health issue is crucial for achieving global commitments expressed in both the ICPD Programme of Action and the Millennium Development Goals. Governments and policymakers must hold themselves accountable to following through on those commitments.

### **References**

- Boland, Reed and Katzive, Laura. 2008. Developments in laws on induced abortion: 1998–2007. *International Family Planning Perspectives*, 34(3): 110-120; <http://www.guttmacher.org/pubs/journals/3411008.html>
- Center for Reproductive Rights (CRR). 2007. *Abortion worldwide: 12 years of reform*. New York: CRR. [http://reproductiverights.org/sites/crr.civicaactions.net/files/documents/pub\\_bp\\_abortionlaws10.pdf](http://reproductiverights.org/sites/crr.civicaactions.net/files/documents/pub_bp_abortionlaws10.pdf)
- Cleland J, Bernstein S, Ezeh A, Faundes A, Glasier A, Innis J. 2006. Family planning: the unfinished agenda. *Lancet*; 368:1810-1827.
- FIGO (International Federation of Gynecology and Obstetrics). 2006. *Ethical issues in obstetrics and gynecology*. London: FIGO. <http://www.figo.org/files/figo-corp/docs/Ethics%20Guidelines%20-%20English%20version%202006%20-2009.pdf>
- International Planned Parenthood Federation (IPPF). 2006. *International Medical Advisory Panel (IMAP) statement on safe abortion*. London: IPPF. <http://www.ippfen.org/NR/rdonlyres/981BE0B4-B04C-4DC4-BD4F-0F7F8339B08F/0/IM-APstatementmay2006.pdf>
- United Nations. 1994. *Programme of action of the International Conference on Population and Development*, Cairo, Egypt, 5–13 September 1994. New York: United Nations. <http://www.unfpa.org/icpd/icpd-programme.cfm>
- United Nations. 1999. *Key actions for the further implementation of the Programme of Action of the International Conference on Population and Development*. A/S-21/5/Add.1. New York: United Nations. [http://www.unfpa.org/upload/lib\\_pub\\_file/561\\_filename\\_icpd5-key-04reprint\\_eng.pdf](http://www.unfpa.org/upload/lib_pub_file/561_filename_icpd5-key-04reprint_eng.pdf)
- World Health Organization (WHO). 2003. *Safe abortion: Technical and policy guidance for health systems*. Geneva: WHO. [http://www.who.int/reproductive-health/publications/safe\\_abortion/safe\\_abortion.pdf](http://www.who.int/reproductive-health/publications/safe_abortion/safe_abortion.pdf)
- World Health Organization (WHO). 2007. *Unsafe abortion: global and regional estimates of incidence of unsafe abortion and associated mortality in 2003*. 5th edition. Geneva: WHO. [http://www.who.int/reproductive-health/publications/unsafeabortion\\_2003/ua\\_estimates03.pdf](http://www.who.int/reproductive-health/publications/unsafeabortion_2003/ua_estimates03.pdf)



**Ipas** *Protecting women's health  
Advancing women's reproductive rights*

P.O. Box 5027 • Chapel Hill, NC 27514 USA  
1.919.967.7052 • [ipas@ipas.org](mailto:ipas@ipas.org) • [www.ipas.org](http://www.ipas.org)

Photos: Martin Adler/Panos Pictures, Gary Calton/Panos Pictures

The photographs used in this publication are for illustrative purposes only; they do not imply any particular attitudes, behaviors, or actions on the part of any person who appears in the photographs.

©2009 Ipas.

Produced in the United States of America.

♻️ Printed on recycled paper.

ICPDMDG-E09