

Training Curriculum – Postabortion care at or after 13 weeks gestation ('second trimester')

PAC Quality Assurance activity - logbook recording

PARTICIPANT HANDOUT

Case: A 23 y.o. G3P2 presents at 20 weeks with intrauterine fetal demise. She is healthy and has had no prior uterine surgeries. She receives mifepristone 200mg and returns 36 hours later at which time she is admitted to hospital. She receives 400mcg vaginally of misoprostol at 0800 AM, her second dose 400mcg vaginally at 1100 AM, and her third dose 400mcg vaginally at 200 PM. She expels without complications one hour after the 3rd dose of misoprostol. She received NSAIDs for pain control. Prior to discharge, she chooses an implant for contraception and this is immediately placed.

Provider Case Logbook for medical management of PAC and/or indicated abortion at or after 13 weeks

Facility name _____ Region _____

Provider name _____ Title/position _____ Address _____

ID	Date	Age	Gravidity	GA in weeks	Indication	Medication doses (track each dose and time)		Expulsion interval in hours	Type of Analgesia	*SAE (Y/N)	Contraception	Other Remarks
						Mifepristone	Misoprostol					

* SAE – Severe Adverse Event: includes the following (bleeding requiring a blood transfusion, infection requiring intravenous antibiotics and/or hospital admission, unintended intra-abdominal surgery, ectopic pregnancy unrecognized at time of procedure, perforation, anesthesia related complication requiring hospitalization or causing seizures, ongoing pregnancy and woman decided not to terminate)

Y/N = Yes /No

Remark: please note here if assistance (manual, MVA, forceps) was needed to remove the placenta.