

## Training Curriculum - Medical abortion at or after 13 weeks gestation ('second trimester')

### Quality Assurance activity - logbook recording

#### PARTICIPANT HANDOUT

A 23 y.o. G3P2 presents at 20 weeks with an undesired pregnancy due to rape. She is healthy and has had no prior uterine surgeries. She receives mifepristone 200mg and returns 36 hours later at which time she is admitted to hospital. She receives 400mcg vaginally of misoprostol at 0800 AM, her second dose 400mcg vaginally at 1100 AM, and her third dose 400mcg vaginally at 200PM. She expels without complications 1 hour after the 3rd dose of misoprostol. She received diclofenac for pain control. Prior to discharge, she chooses a contraceptive implant for contraception.

Provider Case Log Book for Medical Induction of Abortion at or after 13 weeks

Facility name \_\_\_\_\_ Region \_\_\_\_\_

Provider name \_\_\_\_\_ Title/position \_\_\_\_\_ Address \_\_\_\_\_

ID	Date	Age	Gravidity	GA in weeks	Indication	Medication doses (track each dose and time)		Expulsion interval in hours	Type of Analgesia	*SAE (Y/N)	Contraception	Other Remarks
						Mifepristone	Misoprostol					

\* SAE – Severe Adverse Event: includes the following (bleeding requiring a blood transfusion, infection requiring intravenous antibiotics and/or hospital admission, unintended intra-abdominal surgery, ectopic pregnancy unrecognized at time of procedure, perforation, anesthesia related complication requiring hospitalization or causing seizures, ongoing pregnancy and woman decided not to terminate)

Y/N = Yes /No

Remark: please note here if assistance (manual, MVA , forceps) was needed to remove the placenta.